

**4229 FIRST AID KIT - 4229 first aid kit**  
**Honeywell Safety Products USA, INC**

*Disclaimer: This drug has not been found by FDA to be safe and effective, and this labeling has not been approved by FDA. For further information about unapproved drugs, [click here](#).*

-----

**4229 First Aid Kit ( alcohol wipes, ammonia, triple, aypanal, Pain stoppers- 68140FEDX)**

**Ammonia Inhalent**  
***Active ingredient***

Ammonia 15%

**Ammonia Inhalent**  
***Purpose***

Respiratory stimulant

**Ammonia Inhalent**  
***Uses***

- to prevent or treat fainting

**Ammonia Inhalent**  
***Warnings***

**For external use only**

**Do not use**

- if you have breathing problems such as asthma or emphysema

**Stop use and ask a doctor if**

- condition persists

**Keep out of reach of children**

If swallowed get medical help or contact a Poison Control Center right away.

**Ammonia Inhalent**  
***Directions***

- hold inhalant away from face and crush ampoule between thumb and forefinger at position indicated on sleeve.
- hold near nostrils for inhalation of volatile vapor

**Ammonia Inhalent**  
***Other information***

- store at room temperature away from light

**Ammonia Inhalent**  
***Other information***

- store at room temperature away from light

**Ammonia Inhalent*****Inactive ingredients***

alcohol USP, FD&C red #40, lavender oil, lemon oil fcc, nutmeg oil, purified water

**Ammonia Inhalent*****Questions or Comments***

1-800-430-5490

**Alcohol Wipe*****Active ingredient***

Isopropyl alcohol 70%

**Alcohol Wipe*****Purpose***

First aid antiseptic

**Alcohol Wipe*****Uses***

- first aid to help prevent infection in minor cuts, scrapes, and burns

**Alcohol Wipe*****Warnings*****For external use only****Do not use**

- in the eyes
- over large areas of the body

**Ask a doctor before use if you have**

- deep or puncture wounds
- animal bites
- serious burn

**When using this product**

- do not use longer than one week unless directed by a doctor

**Stop use and consult a doctor**

- if condition persists or gets worse

**Keep out of reach of children**

- If swallowed, get medical help or contact a Poison Control Center right away.

**Alcohol Wipe**

**Directions**

- clean the affected area
- apply wipe to affected area 1 to 3 times daily
- may be covered with a sterile bandage
- discard wipe after single use

**Alcohol Wipe****Other information**

store at room temperature 15<sup>0</sup> to 25<sup>0</sup> C (59<sup>0</sup> to 77<sup>0</sup>F)

**Alcohol Wipe****Inactive ingredient**

water

**Alcohol Wipe****Questions**

1-800-430-5490

**Aypanal****Active ingredient**

Acetaminophen 325 mg

**Aypanal****Purpose**

Pain reliever/ fever reducer

**Aypanaly****Uses**

- temporarily relieves minor aches and pains due to the common cold and headache - temporarily reduces fever

**Keep out of reach of children.**

Keep out of reach of children.

**Aypanal****Warnings**

**Liver warning:** This product contains acetaminophen. Severe liver damage may occur if you take:

- more than 4,000 mg in 24 hours, which is the maximum daily amount
- child takes more than 5 doses in 24 hours, which is the maximum daily amount
- taken with other drugs containing acetaminophen
- adult has 3 or more alcoholic drinks every day while using this product:

**Allergy alert:** Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters

- rash
- If a skin rash occurs, stop use and seek medical help right away.

**Do not use**

- with any other drug containing acetaminophen (prescription or nonprescription).
- If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.

**Ask a doctor before use if you have**

- liver disease

**Ask a doctor or pharmacist before use if**

- you are taking the blood thinning drug warfarin

**Stop using and ask a doctor if**

- pain gets worse or lasts more than 10 days in adults
- pain gets worse or lasts more than 5 days in children under 12 years
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur

**If pregnant or breast-feeding**

If pregnant or breast-feeding, ask a health professional before use.

**Overdose warning**

- In case of accidental overdose, get medical help or contact a Poison Control Center right away.
- Prompt medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

**Aypanal*****Directions*****do not take more than directed (see overdose warning)****▣adults and children 12 years of age or older**

- take two tablets every 4-6 hours while symptoms last
- do not take more than 12 tablets in 24 hours

**children 6 to under 12 years of age**

- take 1 tablet every 4-6 hours while symptoms last
- do not take more than 5 tablets in 24 hours

**children under 6 years consult a doctor****Aypanal*****Other information***

- store at room temperature 15<sup>0</sup> to 30<sup>0</sup> C (59<sup>0</sup> - 86<sup>0</sup> F)
- TAMPER EVIDENT PACKETS- DO NOT USE IF OPEN OR TORN

**Aypanal**

***Inactive ingredients***

corn starch, microcrystalline cellulose, povidone, sodium starch glycolate, stearic acid

**Aypanal*****Questions***

1-800-430-5490

**Triple*****Active ingredient***

Bacitracin zinc 400 units

Neomycin sulfate (5 mg equivalent to 3.5 mg Neomycin base)

Polymyxin B sulfate 5000 units

**Triple*****Purpose***

First aid antibiotic

First aid antibiotic -

First aid antibiotic

**Triple*****Uses***

first aid to help prevent infection in

- minor cuts
- scrapes -
- burns

**Triple*****Warnings***

**For external use only**

**Allergy alert** do not use if you are allergic to any of the ingredients

**Do not use**

- in the eyes
- over large areas of the body

**Stop use and ask a doctor if**

- the condition persists or gets worse
- a rash or other allergic reaction develops
- you need to use longer than 1 week

**Ask a doctor before use if you have**

- a deep or puncture wounds
- animal bites
- serious burns

## **Keep out of the reach of children**

If swallowed, get medical help or contact a Poison Control Center right away.

## **Triple**

### ***Directions***

- clean the affected area
- apply a small amount of the product (an amount equal to the surface area of the tip of a finger) on the area 1 to 3 times daily
- may be covered with a sterile bandage

## **Triple**

### ***Other information***

- store at 15 o to 25 oC (59 o to 77 o F)
- tamper evident sealed packets
- do not use if packet is torn or opened

## **Triple**

### ***Inactive ingredient***

petrolatum

## **Pain Stoppers**

### ***Active ingredient***

**Acetaminophen 110mg**

**Aspirin 162mg (NSAID)\***

**Caffeine 32.4mg**

**Salicylamide 152mg (NSAID)\***

**\*nonsteroidal anti-inflammatory drug**

## **Pain Stoppers**

### ***Purpose***

**Pain reliever/fever reducer**

**Pain reliever/fever reducer**

Diuretic

**Pain reliever/fever reducer**

## **Pain Stoppers**

### ***Uses***

for the temporary relief of minor aches and pains due to:

- common cold
- headache
- muscular aches
- premenstrual and menstrual cramps

## **Pain Stoppers**

### **Warnings**

**Reye's syndrome:** Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

**Allergy alert:** Aspirin may cause a severe allergic reaction which may include:

- hives
- facial swelling
- asthma (wheezing)
- shock

**Allergy alert:** Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash

If skin reaction occurs, stop use and seek medical help right away.

**Liver warning:** This product contains acetaminophen. Severe liver damage may occur if you take:

more than 12 tablets in 24 hours, which is the maximum daily amount  
with other drugs containing acetaminophen

3 or more alcoholic drinks every day while using this product

**Stomach bleeding warning:** This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you:

- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen or others)
- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed

### **Do not use**

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist
- for more than 10 days for pain unless directed by a doctor
- for more than 3 days for fever unless directed by a doctor

### **Ask a doctor before use if**

- stomach bleeding warning applies to you
- you have a history of stomach problems, such as heartburn
- you are taking a diuretic
- you have high blood pressure, heart disease, liver cirrhosis, or kidney disease

### **Stop use and ask a doctor if**

- symptoms do not improve
- new symptoms occur
- pain or fever persists or gets worse

- you experience any of the following signs of stomach bleeding:
- feel faint
- vomit blood
- have bloody or black stools
- have stomach pain that does not get better
- if ringing in the ears or a loss of hearing occurs, consult a doctor before taking any more of this product.

### **If pregnant or breast-feeding**

- ask a healthcare professional before use.

It is especially important not to use aspirin during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during pregnancy.

### **Keep out of reach of children**

Overdose warning : **In case of accidental overdose, get medical help or contact a Poison Control Center right away. Prompt medical attention is critical for adults as well as children even if you do not notice any signs or symptoms.**

### **Pain Stoppers**

#### ***Directions***

- adults and children 12 years of age and over, take 2 tablets every 4 hours while symptoms persist
- do not take more than 12 tablets in 24 hours
- children under 12 years: consult a doctor

### **Pain Stoppers**

#### ***Other information***

store at a controlled room temperature 15<sup>0</sup> -30<sup>0</sup> C (59<sup>0</sup> -86<sup>0</sup> F)  
TAMPER EVIDENT-DO NOT USE IF OPEN OR TORN

### **Pain Stoppers**

#### ***Inactive ingredients***

FD&C Yellow #6, magnesium stearate, microcrystalline cellulose, povidone, starch, stearic acid,

### **Pain Stoppers**

#### ***Questions or Comments?***

1-800-430-5490

### **4229**

#### **68140FEDX Kit Contents**

1 WOVEN 2" X 3" 25/BOX

1 FINGERTIP "T" WOVEN 40/BOX

1 1X3 WOVEN SING 50/BOX



1 SWIFT KNUCKLE 40/BX

1 AMMONIA INHALANTS 10 PER

1 NITRILE GLOVES 2PR BBP  
1 ADHESIVE TAPE W/P 1" X 10YDS  
1 GAUZE BANDAGE 2"X2 YDS STRETCH GZ  
1 FIRST AID GUIDE ASHI  
1 BLOODSTOPPER  
1 CTA STRL 6" SGL TIP 12/2  
1 ALCOHOL WIPES 50'S  
1 AYPANAL NON-ASP IND 2/ENV 100  
1 PAIN STOPPERS IND PK 2ENV 250  
1 TRIPLE BIOTIC .5 GRAM PKT 20  
1 SCISSOR BDGE 4" RED PLS HDL  
1 KIT TWEEZER 3 1/2" SLANTED  
1 F A KIT EMPTY BLANK 140  
LBL STOCK 6-3/8"X4"  
1 LBL STOCK 3"x1-7/8"  
1 TRI BNDG NON WOVEN 40"X40"X56"

**Ammonia inhalent**  
***Principal Display Panel***


796006 Rev. E Unit Carton Printing Plate for "A" size carton.




796006 Rev. E (page 3 of 3)

# Principal Display Panel

825362T Rev. A  
 white printed  
 Pms 072 blue, Pms 186 red,  
 Pms 7459 blue and black



# Aypanal Principal Display Panel



# Triple Principal Display Panel



# Pain Stoppers Principal Display Panel





Distributed by Honeywell Safety Products USA, Inc Smithfield, RI 02917

## 4229 FIRST AID KIT

4229 first aid kit kit

### Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:0498-4229
--------------	----------------	--------------------	---------------

**Packaging**

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:0498-4229-01	1 in 1 KIT; Type 0: Not a Combination Product	10/18/2018	

**Quantity of Parts**

Part #	Package Quantity	Total Product Quantity
Part 1	20 PACKET	10 g
Part 2	125 PACKET	250
Part 3	10 AMPULE	3 mL
Part 4	50 POUCH	20 mL
Part 5	50 PACKET	100

**Part 1 of 5****TRIPLE ANTIBIOTIC**

bacitracin zinc, polymyxin b sulfate, neomycin sulfate ointment

**Product Information**

Item Code (Source)	NDC:0498-0750
Route of Administration	TOPICAL

**Active Ingredient/Active Moiety**

Ingredient Name	Basis of Strength	Strength
POLYMYXIN B SULFATE (UNII: 19371312D4) (POLYMYXIN B - UNII:J2VZ07J96K)	POLYMYXIN B	5000 [iU] in 1 g
NEOMYCIN SULFATE (UNII: 057Y626693) (NEOMYCIN - UNII:I16QD7X297)	NEOMYCIN	3.5 mg in 1 g
BACITRACIN ZINC (UNII: 89Y4M234ES) (BACITRACIN - UNII:58H6RWO52I)	BACITRACIN	400 [iU] in 1 g

**Inactive Ingredients**

Ingredient Name	Strength
PETROLATUM (UNII: 4T6H12BN9U)	

**Product Characteristics**

Color	white	Score	
Shape		Size	
Flavor		Imprint Code	
Contains			

**Packaging**

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:0498-0750-36	0.5 g in 1 PACKET; Type 0: Not a Combination Product		

## Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC monograph final	part333B	09/19/2018	

## Part 2 of 5

### PAIN STOPPERS

acetaminophen, caffeine, aspirin, salicylamide tablet

## Product Information

Item Code (Source)	NDC:0498-2422
Route of Administration	ORAL

## Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
ACETAMINOPHEN (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	110 mg
ASPIRIN (UNII: R16CO5Y76E) (ASPIRIN - UNII:R16CO5Y76E)	ASPIRIN	162 mg
CAFFEINE (UNII: 3G6A5W338E) (CAFFEINE - UNII:3G6A5W338E)	CAFFEINE	32.4 mg
SALICYLAMIDE (UNII: EM8BM710ZC) (SALICYLAMIDE - UNII:EM8BM710ZC)	SALICYLAMIDE	152 mg

## Inactive Ingredients

Ingredient Name	Strength
STEARIC ACID (UNII: 4ELV7Z65AP)	
MAGNESIUM STEARATE (UNII: 70097M6I30)	
POVIDONE K30 (UNII: U725QWY32X)	
CELLULOSE, MICROCRYSTALLINE (UNII: OP1R32D61U)	
FD&C YELLOW NO. 6 (UNII: H77VEI93A8)	

## Product Characteristics

Color	orange (Bright Orange)	Score	no score
Shape	ROUND	Size	11mm
Flavor		Imprint Code	FR;2
Contains			

## Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:0498-2422-01	2 in 1 PACKET; Type 0: Not a Combination Product		

## Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC monograph not final	part343	01/02/2017	

## Part 3 of 5

### AMMONIA INHALENT

ammonia inhalent inhalant

## Product Information

Item Code (Source)	NDC:0498-3334
Route of Administration	RESPIRATORY (INHALATION)

## Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
AMMONIA (UNII: 5138Q19F1X) (AMMONIA - UNII:5138Q19F1X)	AMMONIA	0.045 g in 0.3 mL

## Inactive Ingredients

Ingredient Name	Strength
ALCOHOL (UNII: 3K9958V90M)	

## Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:0498-3334-00	0.3 mL in 1 AMPULE; Type 0: Not a Combination Product		

## Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
unapproved drug other		09/18/2018	

## Part 4 of 5

### ALCOHOL WIPE

isopropyl alcohol swab

## Product Information

Item Code (Source)	NDC:0498-0143
--------------------	---------------

Route of Administration TOPICAL

### Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
ISOPROPYL ALCOHOL (UNII: ND2M416302) (ISOPROPYL ALCOHOL - UNII:ND2M416302)	ISOPROPYL ALCOHOL	0.7 mL in 1 mL

### Inactive Ingredients

Ingredient Name	Strength
WATER (UNII: 059QF0KO0R)	

### Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:0498-0143-04	0.4 mL in 1 POUCH; Type 0: Not a Combination Product		

### Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC monograph not final	part333A	09/18/2018	

## Part 5 of 5

### AYPANAL NON-ASPIRIN

acetaminophen tablet

### Product Information

Item Code (Source)	NDC:0498-2001
Route of Administration	ORAL

### Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
ACETAMINOPHEN (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	325 mg

### Inactive Ingredients

Ingredient Name	Strength
STARCH, CORN (UNII: O8232NY3SJ)	
STEARIC ACID (UNII: 4ELV7Z65AP)	
POVIDONE (UNII: FZ989GH94E)	
SODIUM STARCH GLYCOLATE TYPE A CORN (UNII: AG9B65PV6B)	

## Product Characteristics

Color	white	Score	2 pieces
Shape	ROUND	Size	10mm
Flavor		Imprint Code	circle;U
Contains			

## Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1		2 in 1 PACKET; Type 0: Not a Combination Product		

## Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC monograph not final	part343	04/10/2012	

## Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
unapproved drug other		10/18/2018	

**Labeler** - Honeywell Safety Products USA, INC (079287321)

## Establishment

Name	Address	ID/FEI	Business Operations
James Alexander		040756421	manufacture(0498-3334)

## Establishment

Name	Address	ID/FEI	Business Operations
Honeywell Safety Products USA, INC		079287321	pack(0498-4229)

## Establishment

Name	Address	ID/FEI	Business Operations
Ultra Seal Corporation		085752004	manufacture(0498-2001, 0498-2422)

## Establishment

Name	Address	ID/FEI	Business Operations
Water-Jel Technologies		155522589	manufacture(0498-0750)

## Establishment

Name	Address	ID/FEI	Business Operations
Changzhou Maokang Medical		421317073	manufacture(0498-0143)

